



APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: METHOD FOR IDENTIFYING MICROBIAL PROLIFERATION GENES

**APPLICANT: JUDITH ZYSKIND, a United States Citizen
and a resident of La Jolla, California**

and

**R. ALLYN FORSYTH, a United States Citizen
and a resident of San Diego, California.**

**Fish & Richardson P.C.
4350 La Jolla Village Drive, Suite 500
San Diego, California 92122
Telephone: 858/678-5070
Facsimile: 858/678-5099**

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL688321692US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

Date of Deposit March 13, 2001

Signature

Gildardo Vargas, Jr.

Typed or Printed Name of Person Signing Certificate

Sub
B1
SCANNED, #

SUB
B1
IUS
B2

METHOD FOR IDENTIFYING MICROBIAL PROLIFERATION GENES

Statement as to Federally Sponsored Research

5 Research reported herein was supported in part by the National Science Foundation and the National Institutes of Health. The United States government may have certain rights in this invention.

Background of the Invention

15 Genes that encode proteins essential for microbial growth or viability are useful targets for antibiotics because inhibition of such proteins by an antibiotic can reduce or eliminate the spread of infectious disease in a patient. The emergence of bacteria resistant to multiple antibiotics has led to renewed interest in isolating variants of known antibiotics and also in identifying new required genes and the corresponding gene products that could serve as new targets for novel antibiotics.

20 Approximately fifteen different bacterial proteins encoded by essential genes have been used as targets for antibiotics. Such target proteins include ribosomal proteins, gyrase, RNA polymerase and proteins involved in the synthesis of the peptidoglycan layer and its precursors. However, there are estimated to be more than 4000 putative genes in the genome of the bacterium *Escherichia coli* and it is not known how many of these genes are required for growth and/or viability. The discovery of additional required genes could facilitate the search for new

antibiotics.

Microbial genes required or essential for growth and/or viability have been identified by two major techniques. One approach is to determine the nucleotide sequence of the genome of a microbial species of interest. Sequence information is then compared to sequences in computer databases to identify possible functions for the putative gene sequences. To prove that a putative gene identified only by sequence comparisons is in fact an essential gene, however, a null or "knockout" mutation must be made in the gene. It must be shown that a microorganism containing the null mutation cannot survive unless it is complemented by a wild type allele of the gene. Such an approach is time consuming and, in some species, can be difficult.

A second approach is to isolate conditionally lethal mutants by means such as chemical mutagenesis. A common type of conditional lethal mutation is a temperature-sensitive mutation, in which the mutant is non-viable at higher temperature such as 42° C but is viable at a lower temperature such as 30° C. The mutation is then mapped and identified by cloning and genetic complementation. The mutagenesis techniques employed are sometimes not completely random, thus reducing the likelihood of identifying a required gene that does not have a "hot spot" for the mutagen used. Moreover, products of some required genes may not be amenable to the formation of conditional lethal mutants. Such essential genes would not be identified by this approach.

Summary of the Invention

The present invention is based on the discovery that microbial genes and their encoded proteins essential for growth and/or viability can be identified by introduction into a

microorganism of an exogenous nucleic acid having sequence similarity to the microbial gene. The exogenous nucleic acid can be, for example, an antisense RNA that is derived from a chromosomal fragment cloned downstream of an inducible promoter. The endogenous gene can be identified as required by the inhibition of proliferation of microorganisms containing or
5 expressing the exogenous nucleic acid, compared to the proliferation of microorganisms that do not contain or are not expressing the nucleic acid. The invention facilitates the rapid, efficient identification of microbial genes not previously known to be required for growth or viability.

It will be evident to those skilled in the art that the invention can be applied to any organism for which it is feasible to 1) isolate and fragment the chromosome of the organism, 2) re-introduce exogenous nucleic acid fragments of the organism or fragments substantially similar to those of the organism back into the organism and 3) culture the organism without the exogenously added nucleic acids.

The present invention advantageously permits the rapid, inexpensive identification of microbial genes that heretofore have not been identified as required for survival. The method comprises the steps of introducing an exogenous nucleic acid into a microorganism and determining the proliferation of the microorganism relative to the proliferation of the microorganism when the exogenous nucleic acid is not present or not expressed. The exogenous nucleic acid has substantial sequence identity to an endogenous microbial gene. The exogenous
20 nucleic acid can be operably linked to a regulatory element effective for controlling expression of the nucleic acid, either in sense orientation or in antisense orientation to the regulatory element. The regulatory element can control expression via chemical induction of the exogenous nucleic acid. The microorganism can be, for example, a bacterium, either gram-negative or gram-

positive, a fungus, or an Archaeobacteria.

The novel genes and the proteins encoded by such genes are promising targets for new antimicrobial agents. A method for identifying such agents comprises the steps of contacting an agent with a microorganism having a proliferation gene and determining the effect of the agent on the proliferation gene or a product of the proliferation gene. The proliferation gene can be endogenous to the microorganism, or can be from another source microorganism. The effect of the agent on the proliferation gene can be, for example, on transcription or on translation of the proliferation gene.

Specific genes and their mRNAs identified by the novel method can be targets for antisense polynucleotide inhibitors, and these antisense polynucleotides can have therapeutic value. A method for inhibiting expression of a microbial proliferation gene comprises contacting a microorganism having a proliferation gene with an antisense polynucleotide having substantial sequence identity to the proliferation gene.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

Brief Description of the Drawings

~~Figure 1 shows the nucleotide sequence of an antisense RNA to *E. coli lepB* mRNA,~~

expressed from the DNA insert in plasmid pJB37.

Figure 2 shows the nucleotide sequence of an antisense RNA to *E. coli viaA* mRNA, expressed from the DNA insert in plasmid pJB53.

Figure 3 shows the nucleotide sequence of a 714 nucleotide *Pst*I-*Hind*III fragment of the *E. coli viaA* gene cloned in pJB53.

Figure 4 shows the nucleotide sequence of an antisense RNA to *E. coli ddlB* mRNA, expressed from the DNA insert in plasmid pJB57.

Figure 5 shows the nucleotide sequence of an antisense RNA to *orf1* of the *E. coli ampG* operon, expressed from the DNA insert in plasmid pJB59.

Figure 6 is a bar graph showing the effect of inducing *lepB*, *ddlB*, *viaA* or *orf1* antisense RNA expression on colony formation in *E. coli* DH5 α . Striped bars indicate colony forming units (CFUs) per ml of culture for strains grown on semi-solid media in the absence of the inducer IPTG. Solid bars indicate CFUs/ml for strains grown in the presence of IPTG.

Figure 7 is a graph showing growth curves of *E. coli* DH5 α containing pJB37, pJB57 and pJB59 in liquid media in the presence and absence of IPTG.

Figure 8 is a graph showing the viability of DH5 α strains at various times after induction of antisense RNA expression. Strains contained the following plasmids: squares: pJB37; diamonds: pJB53; circles: pJB57; and triangles; pJB59.

Figure 9 is a graph showing the effect of varying concentrations of inducer on the viability of DH5 α containing pJB37 (panel A) or pJB53 (panel B).

Figure 10 is a bar graph comparing the effect of ampicillin or spectinomycin selection on viability of DH5 α strains expressing *lepB*, *ddlB*, or *orf1* antisense RNAs. Ampicillin selection

(narrow stripes), ampicillin selection with 1 mM IPTG (gray), spectinomycin selection (wide stripes) and spectinomycin selection with 1 mM IPTG (solid).

5 ~~Figure 11 shows the nucleotide sequence of an 836 nucleotide *Pst*I/*Hind*III fragment of the *E. coli secA* gene cloned in pJB3.~~

a Figure 12 shows the nucleotide sequence of an antisense RNA complementary to *E. coli secA* mRNA.

Description of the Preferred Embodiments

10 The present invention provides an efficient strategy for identifying microbial proliferation genes. Proliferation genes are genes that are necessary for proliferation, *i.e.*, in the absence of gene transcript and/or gene product, growth or viability of the microorganism is reduced or eliminated.

15 Microorganisms can be used in the invention as a pure culture or may be used as a mixed culture, *i.e.*, more than one strain or species may be used in the novel method. Microorganisms suitable for use in the method include both prokaryotes and eukaryotes. Bacterial species suitable for use in the method include substantially all bacterial species, both animal and plant pathogenic species, as well as non-pathogenic species. Preferred species are pathogens of mammals, especially pathogens of humans such as *Staphylococcus aureus*.

20 Either gram-negative and gram-positive bacterial species can be used, *e.g.*, gram-positive *Enterococcus*, *Bacteriodes*, *Clostridium* and *Staphylococcus* species and gram-negative *Escherichia*, *Pseudomonas*, *Haemophilus*, *Enterobacter*, *Vibrio*, *Salmonella*, *Helicobacter* and *Moraxella* species. *Mycobacterium* species such as *Mycobacterium tuberculosis* are also suitable

for use in the method disclosed herein. Eukaryotic microbes suitable for use in the method include, without limitation, *Cryptococcus neoformans*, *Nocardia spp.*, *Coccidioides immitis*, *Toxoplasma spp.*, *Leishmania spp.*, *Pneumocystis*, *Trypanosoma spp.*, *Giardia spp.*, *Plasmodia*
5 *spp.* (e.g., *Plasmodium falciparum*), *Histoplasma capsulatum*, and *Candida albicans*.

The novel method for identifying microbial proliferation genes comprises the step of introducing an exogenous nucleic acid into a microorganism. The exogenous nucleic acid may be obtained, for example, from a restriction digest of total genomic (e.g., chromosomal and episomal) DNA from a microorganism of interest, or from total genomic DNA that has been subjected to physical shear. Sheared DNA may result in a more random set of fragments than would otherwise be generated by restriction digestion. Nucleic acid fragments obtained by partial or total restriction digestion or by shearing can be size selected by agarose gel electrophoresis or sucrose gradients, if desired. Exogenous nucleic acid can also be obtained by chemical synthesis (synthetic DNA having a random sequence or synthetic DNA based on information in a nucleic acid sequence database), from a cDNA library, or by other means known to the art.

An exogenous nucleic acid of the present invention can be in the form of RNA or in the form of DNA, including cDNA, synthetic DNA or genomic DNA. The DNA can be double-stranded or single-stranded and, if single-stranded, can be either a coding strand or non-coding
20 strand. RNA can be, for example, mRNA or a combination of ribo- and deoxyribonucleotides.

An exogenous nucleic acid of the present invention has substantial sequence identity to an endogenous microbial proliferation gene. It should be appreciated, however, that it is not necessary to have prior knowledge of sequence identity to an endogenous gene. In fact, an

advantage of the present method is that a plurality of randomly generated nucleic acid fragments, whose identity to particular endogenous genes has not been determined, can be screened to identify a fragment (or fragments) that inhibit microbial proliferation. The relationship between such a fragment and the endogenous microbial gene can then be determined.

An exogenous nucleic acid has at least about 70% sequence identity, preferably at least about 80% sequence identity, more preferably at least about 90% sequence identity to the endogenous microbial proliferation gene. Sequence identity can be determined, for example, by computer programs designed to perform single and multiple sequence alignments. Nucleic acids having at least about 70% nucleotide sequence identity to the endogenous gene can also be identified by their ability to hybridize to the endogenous gene under conditions of moderate stringency. Nucleic acids having at least about 80% sequence identity, or at least about 90% sequence identity, or at least about 95% sequence identity to the endogenous gene can hybridize under high stringency conditions.

Hybridization can be measured by Southern analysis (Southern blotting), a method by which the presence of DNA sequences in a target nucleic acid mixture are identified by hybridization to a labeled oligonucleotide or DNA fragment probe. Southern analysis typically involves electrophoretic separation of DNA digests on agarose gels, denaturation of the DNA after electrophoretic separation, and transfer of the DNA to nitrocellulose, nylon, or another suitable membrane support for analysis with a radiolabeled, biotinylated, or enzyme-labeled nucleic acid as described in sections 9.37-9.52 of Sambrook *et al.*, MOLECULAR CLONING: A Laboratory Manual, Vol. 2, Chapter 11, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989).

An exogenous nucleic acid can hybridize to the endogenous gene under moderate stringency conditions or under high stringency conditions. High stringency conditions are used to identify nucleic acids that have a high degree of homology or sequence identity to one another.

5 High stringency conditions can include the use of a denaturing agent such as formamide during hybridization, *e.g.*, 50% formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, and 75 mM sodium citrate at 42°C. Another example is the use of 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS. Alternatively, low ionic strength and high temperature can be employed for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate (0.1X SSC); 0.1% sodium lauryl sulfate (SDS) at 65°C.

Moderate stringency conditions are hybridization conditions used to identify nucleic acids that have less homology or identity to one another than do nucleic acids identified under high stringency conditions. Moderate stringency conditions can include the use of higher ionic strength and/or lower temperatures for washing of the hybridization membrane, compared to the ionic strength and temperatures used for high stringency hybridization. For example, a wash solution comprising 0.060 M NaCl/0.0060 M sodium citrate (4X SSC) and 0.1% sodium lauryl sulfate (SDS) can be used at 50°C, with a last wash in 1X SSC, at 65°C. Alternatively, a 20 hybridization wash in 1X SSC at 37°C can be used.

Hybridization can also be done by Northern analysis (Northern blotting), a method used to identify RNAs that hybridize to the nucleic acid used as the probe. The probe is labeled with a

radioisotope such as ^{32}P , by biotinylation, or with an enzyme. The RNA to be analyzed can be electrophoretically separated on an agarose or polyacrylamide gel, transferred to nitrocellulose, nylon, or other suitable membrane, and hybridized with the probe, using standard techniques well known in the art such as those described in sections 7.39-7.52 of Sambrook *et al.*, *supra*.

The exogenous nucleic acid can be less than about 400 nucleotides in length, *e.g.*, from about 10 to about 400 nucleotides in length. An exogenous nucleic acid can also be greater than about 400 nucleotides in length, *e.g.*, from about 10 to about 5,000 nucleotides in length. It is generally preferred that an exogenous nucleic acid of about 15 to about 1,500 nucleotides in length be used in the method, *e.g.*, at least about 15 nucleotides, or at least about 30 nucleotides, or at least about 100 nucleotides.

In some embodiments, the nucleotide sequence of the exogenous nucleic acid has substantial complementarity to the coding sequence of the endogenous microbial proliferation gene, as in an antisense RNA. In other embodiments, the nucleotide sequence of the exogenous nucleic acid is the same as or substantially similar to the coding strand sequence of the endogenous gene, *e.g.*, is a sense RNA. An exogenous nucleic acid preferably exerts a microbiostatic (*e.g.*, bacteriostatic or fungistatic) or microbiocidal (*e.g.*, bacteriocidal or fungicidal) effect by antisense inhibition of a microbial proliferation gene. For example, an exogenous nucleic acid can be DNA that is transcribed into an RNA molecule complementary (antisense) to mRNA produced by the endogenous microbial gene. In such an embodiment, the exogenous nucleic acid is the DNA. The expressed antisense RNA has the capability to inhibit translation of the mRNA and thereby inhibit the proliferation of the microbial cells.

There are several possible modes of action for antisense molecules. For example, the

Shine Dalgarno sequence and start codon of a bacterial proliferation gene mRNA could be targeted by an antisense nucleic acid, and thereby inhibit ribosomal loading onto the mRNA. Alternatively, a hybrid molecule could be formed between an antisense nucleic acid and a portion of the coding region of the microbial proliferation gene mRNA. In this case, the mechanism of antisense inhibition could be premature translational termination, due to inhibition of ribosome movement or decreased mRNA stability. As another alternative, a hybrid could be formed between an antisense nucleic acid and a portion of the microbial proliferation gene mRNA near the transcriptional terminator. In addition, a triple helix could form between the proliferation gene and the exogenous nucleic acid, thereby inhibiting transcription of the proliferation gene. In some instances, it is possible that more than one of the above mechanisms, or a different mechanism, may be responsible for the effect on proliferation. The present invention encompasses these and other mechanisms.

Alternatively, an exogenous nucleic acid exerts a microbiostatic or microbiocidal effect by sense inhibition of a microbial proliferation gene, *i.e.*, dominant negative suppression. For example, an exogenous nucleic acid can be DNA that is transcribed into an RNA molecule having a sequence substantially identical to mRNA produced by the endogenous microbial gene. The expressed sense RNA has the capability to inhibit the proliferation of the microbial cells, possibly by producing proteins or truncated peptides that interfere with normal function of the endogenous proliferation gene product.

As discussed herein, it is preferable to obtain a plurality of exogenous nucleic acids, *e.g.*, a library of restriction fragments of a microbial genome. Such a population of exogenous nucleic acids has substantial sequence identity to a large number of endogenous microbial genes,

preferably the entire genome. The population is screened in parallel for an effect on microbial proliferation; those fragments exerting an inhibitory effect on growth and/or viability are selected for further study.

5 Typically, an exogenous nucleic acid is introduced into a microorganism on a vector, *e.g.*, a plasmid or phage vector. Illustrative examples of vectors are the pLEX series of plasmids described in Diederich, L. *et al.*, *Biotechniques* 16:916-923 (1994). Other suitable vectors are known in the art. If desired, assays can be used to confirm that the exogenous nucleic acid has been effectively introduced in order to confirm that the antiproliferative effect is due to the exogenous nucleic acid and not due to some other factor.

10 Techniques are known in the art for the introduction of exogenous nucleic acid into microorganisms, *e.g.*, gram-negative as well as gram-positive bacteria. Techniques for introducing exogenous nucleic acid include, without limitation, lambda phage *in vitro* packaging, freeze-thaw mediated transformation, CaCl_2 -mediated transformation, liposomes, electroporation, natural transformation, conjugation and particle gun transformation.

15 An exogenous nucleic acid preferably is operably linked to at least one suitable regulatory sequence to facilitate expression of the exogenous nucleic acid. Regulatory sequences typically do not themselves code for a gene product. Instead, regulatory sequences control the expression level of the exogenous nucleic acid. An exogenous nucleic acid can be operably
20 linked to a suitable inducible regulatory sequence in antisense orientation or in sense orientation.

 Examples of regulatory sequences are known in the art and include, without limitation, promoters induced in response to chemical inducers, promoters induced in response to environmental changes (temperature shifts, radiation and the like) and promoters repressed in

response to the presence of chemical agents. Examples of promoters induced in the presence of chemical inducers include the *E. coli lac* promoter (induced with isopropyl- β -D-thiogalactopyranoside, also known as IPTG), the *tet* promoter (induced with tetracycline) and the hybrid *tac* promoter (induced with IPTG). Examples of promoters repressed in response to a chemical repressor include the *trp* promoter (repressed by tryptophan, a corepressor). Other suitable regulatory sequences may be included if desired. For example, the *rrnB* transcriptional terminator 3' non-coding region, *rrnBt1t2* may be used.

Molecular biology techniques for linking an exogenous nucleic acid to a regulatory sequence effective for controlling expression of the nucleic acid are known in the art. In some embodiments, more than one regulatory element is operably linked to the exogenous nucleic acid, provided that the nucleic acid can be expressed when desired.

The novel method also includes the step of determining the effect of the exogenous nucleic acid on microbial proliferation, relative to the effect in the absence of the exogenous nucleic acid. For example, the exogenous nucleic acid can be operably linked to an inducible promoter and the effect on proliferation can be measured after induction of expression. The effect on proliferation is compared to microbial proliferation when the nucleic acid is not induced and expressed. Means for determining the effect on proliferation include, for example, measuring colony-forming units (CFU), measuring light scattering (optical density), measuring the number of microorganisms in a particle counter, measuring respiratory activity, measuring fluorescence of cell cultures or individual cells after addition of fluorescent dyes and measuring incorporation of precursors to macromolecular (RNA, DNA, protein, cell wall) synthesis, measuring uptake of metabolites and measuring transport.

Typically, microbial cells are cultured for a period of time to allow the exogenous nucleic acid to inhibit endogenous gene expression. Culturing may occur on solid media or in liquid media, on complex media or minimal media. Culturing may be carried out during an exponential growth phase, stationary growth phase, or during a stress phase. Nevertheless, it should be appreciated that the effect on proliferation can be determined without culturing the microorganism. For example, inhibition of a proliferation gene can be identified by a rapid effect on respiratory activity, a measurement that can be carried out under conditions that do not support long-term growth and viability. Techniques for determining the effect on proliferation that do not involve culturing of microorganisms are within the scope of the present invention.

Inhibition of bacterial growth is readily determined by comparative or replica plating. Bacteria are plated on solid media under conditions in which there is no production or expression of exogenous nucleic acid. Bacterial colonies are subsequently replica plated to a second plate of solid media under conditions in which production or expression of exogenous nucleic acid occurs. Clones that cannot grow in the presence of exogenous nucleic acid are chosen from the first plate for identification.

Culturing may occur for a period of time appropriate for the strain or species being tested. For example, about a 3 hour delay may occur before growth inhibition is observable with an antisense mechanism. Expressed antisense RNA must bind to and inhibit translation of its chromosomally encoded mRNA counterpart. Reduction in protein gene product may be detectable after 1 hour, but even if translational inhibition were 100% it may take several generations to "dilute out" the active gene product below levels required for viability.

It is often preferable to measure the effect on proliferation under culture conditions

similar to those in which it is desired to determine the requirement for the endogenous gene. For example, proliferation genes in a microbial human pathogen are preferably identified under conditions that relate to those found when the pathogen has infected a human, because genes required under such conditions are generally of more interest. It can be useful in some instances to identify proliferation genes under more than one set of conditions, *i.e.*, to determine if the gene is conditionally required. However, the most useful microbial proliferation genes are those genes necessary for growth and/or viability under substantially all conditions, because protein products of such genes are more desirable as targets for novel antibiotics.

Proliferation genes are often highly conserved among different species, for example, different bacterial species. Thus, microbial genes that have been identified as proliferation genes in one species or strain by the method disclosed herein can be used to search for related genes in other species or strains.

In some embodiments, an exogenous nucleic acid is not introduced on a vector (*e.g.*, a plasmid or phage vector), but is introduced as a non-replicating polynucleotide in an amount sufficient to exert an effect on the corresponding endogenous gene without the need for amplification of the amount *in vivo*. Such exogenous nucleic acids typically are antisense polynucleotides, including both short RNA oligonucleotides, typically 10-50 nucleotides in length, as well as longer RNA fragments that may exceed the length of the endogenous microbial gene itself. Antisense nucleic acids useful for the present invention are substantially complementary to regions of the corresponding proliferation gene mRNA. Hybridization of antisense nucleic acids to their target transcripts can be highly specific as a result of complementary base pairing. The capability of antisense nucleic acids to hybridize is affected by

such parameters as length, chemical modification and secondary structure of the transcript which can influence nucleic acid access to the target site. An antisense polynucleotide can also be a DNA segment that synthesizes an antisense RNA inside the cell.

5 In selecting the preferred length for a given nucleic acid, a balance must be struck to gain the most favorable characteristics. Shorter nucleic acids such as 10-to 15-mers, while offering higher cell penetration, have lower gene specificity. In contrast, while longer nucleic acids of 20-30 bases offer better specificity, they show decreased uptake kinetics into cells. See Stein *et al.*, "PHOSPHOROTHIOATE OLIGODEOXYNUCLEOTIDE ANALOGUES" in Oligodeoxynucleotides - Antisense Inhibitors of Gene Expression, Cohen, Ed. McMillan Press, London (1988).

The present invention encompasses both oligomeric nucleic acid moieties of the type found in nature, such as the deoxyribonucleotide and ribonucleotide structures of DNA and RNA. The invention also encompasses synthetic, modified analogues that are capable of binding to an endogenous microbial gene or its mRNA.

In some embodiments, therefore, an antisense polynucleotide of the present invention is a phosphorothioate or methyl phosphonate-linked analogue, which can be nuclease-resistant and thus have increased stability. Alternatively, an antisense polynucleotide can be a peptide nucleic acid, which can be nuclease-resistant and have high affinity for a complementary strand. See,
20 e.g., U.S. Patents 5,539,082 and 5,539,083. Nucleic acids can also have adducts that permit cross-linking between strands of an antisense or sense hybrid. Persons of ordinary skill in this art will be able to select other linkages for use in the invention. These modifications also may be designed to improve the cellular uptake and stability of the nucleic acids.

Phosphorothioate modified DNA nucleic acids typically are synthesized on automated DNA synthesizers available from a variety of manufacturers, *e.g.*, available from Perkin-Elmer/Applied Biosystems (Foster City, CA). These instruments are capable of synthesizing nanomole amounts of nucleic acids as long as 100 nucleotides. Shorter nucleic acids synthesized by modern instruments are often suitable for use without further purification. If necessary, nucleic acids may be purified by polyacrylamide gel electrophoresis or reverse phase chromatography. See Sambrook *et al.*, MOLECULAR CLONING: A Laboratory Manual, Vol. 2, Chapter 11, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989).

Illustrative embodiments of antisense polynucleotides includes polynucleotides having substantial sequence complementarity to the *E.coli lepB*, *viaA*, *ddlB*, *orf1*, and *secA* mRNAs, as well as oligonucleotide subfragments thereof, *e.g.*, about 10 to about 100 nucleotides in length, or about 10 to about 60 nucleotides in length, or about 20 to about 50 nucleotides in length. A desired antisense oligonucleotide can be readily identified by preparing and testing overlapping portions of, *e.g.*, an RNA of Figure 1, 2, 4-5 or 12 for antiproliferative effects as described herein. For example, antisense oligonucleotides about 30 nucleotides in length, each having a 10 nucleotide overlap with adjacent oligonucleotides, can be synthesized and tested for antiproliferative effects. Those oligonucleotides having such an effect are useful as antiproliferative agents as described herein.

Once a microbial proliferation gene has been discovered, the invention provides a method for identifying an antimicrobial agent targeted at the novel gene or gene product. The method includes the step of contacting an agent to be tested for its antimicrobial properties with a microorganism having a proliferation gene. The proliferation gene present in the microorganism

may be a gene endogenous to the microorganism. Alternatively, the proliferation gene can be one that has been introduced from a second microorganism. For example, a proliferation gene from a species for which it is more difficult to evaluate an agent can be introduced into a species for which it is less difficult to evaluate an agent, in order to facilitate testing of various agents.

Contacting includes conditions which allow the test agent to exert an effect, if any, on microbial proliferation, and can occur either in solution or in solid phase. In preferred embodiments, a plurality of agents are tested, *e.g.*, a combinatorial library can be used for screening. The agent is determined to be an antimicrobial agent when the agent is found to exert an effect on a microbial proliferation gene, its mRNA, or its encoded protein as described herein.

The novel screening method is useful for identifying new classes of antimicrobial agents directed at the endogenous microbial proliferation protein targets identified as described herein. Agents suitable for use in the method include inorganic chemical compounds, organic chemical compounds, peptidomimetics, and biologic compounds such as peptides, polypeptides, oligonucleotides and polynucleotides. Typically, the method is useful for identifying an agent that is itself microbiostatic or microbiocidal, *e.g.*, is potentially useful as an antibiotic. In some instances, an agent is useful for identifying a class of structurally related agents that will have efficacious microbiostatic or microbiocidal properties.

If desired, indirect means may be used to determine the effect of an agent on microbial proliferation. For example, an agent can be tested for its effect on enzyme activity if the proliferation gene encodes or is suspected of encoding an enzyme. If a protein does not have an easily assayable enzymatic activity, immunoassays can be carried out with antibody raised against a polypeptide or peptide product of a microbial proliferation gene, in order to identify

effects on transcription and translation of the endogenous gene. Northern blots or *in vitro* transcription assays can be used to determine the effect of the agent on transcription of the microbial proliferation gene. Such assays can be carried out on viable cells, on crude cell lysates
5 or even on purified preparations of the endogenous gene, mRNA or gene product. In addition, structural bioinformatics can be used to identify putative inhibitors and likely mechanisms by which an agent exerts its microbiostatic or microbiocidal effect.

Agents identified by the method can be further evaluated by any method usually applied to the determine the efficacy and pharmacological activity of an antimicrobial agent.

Once a gene has been identified as a proliferation gene, it can be further characterized. Some identified genes will have been previously known as proliferation genes. Others, although having been previously sequenced, will not have been identified as proliferation genes. Others, although having been previously sequenced, are tentatively identified as proliferation genes based on sequence homology to an proliferation gene from another organism. Still others will not have been previously identified or characterized and can be cloned, if desired, using known techniques such as polymerase chain reaction (PCR) technology. See, *e.g.*, U.S. Patent 4,683,195, and subsequent modifications of the procedure described therein. Such genes can also be sequenced by known techniques such as dideoxy sequencing; open reading frames (ORFs) and putative regulatory elements assigned to various regions of the sequenced gene.
20 Homology searches of nucleic acid and protein databases, such as the GenBank, EMBL, and Swiss Prot databases, are useful for deducing possible biochemical and metabolic functions of genes identified as required by the method disclosed herein.

Illustrative examples of proliferation genes include, for example, genes having substantial

sequence identity to the *E. coli viaA* gene, the *E. coli secA* gene, the *E. coli orf1* gene, the *E. coli ddlB* gene and the *E. coli lepB* gene. Such genes can be identified as proliferation genes by sense or antisense inhibition.

5 In some cases, genes are identified as proliferation genes based on the effect observed with sense constructs. For example, genes having substantial sequence identity to the *E. coli ugpB* gene and a gene having homology to the *E. coli fimF* and *fimD* genes can be considered proliferation genes because bacterial growth and/or viability is reduced or eliminated when a sense construct to such genes is expressed. Bacterial inhibition by antisense constructs would not ordinarily be expected to occur for such genes. Nevertheless, it should be appreciated that sense constructs that result in bacterial inhibition can be useful to derive antiproliferative agents. For example, peptides useful as antibiotics can be obtained, based on peptides made by the sense construct. Alternatively, nucleic acids useful as antibiotics can be obtained based on the RNA made by the sense construct.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
1018

Example 1.

Generation of *E. coli* Chromosomal Library

Bacterial strains, plasmids and media. Luria Broth was made as previously described (Sambrook *et al.*, 1989). M9 media (Sambrook *et al.*, 1989) was supplemented with 0.5% casamino acids and 0.2% glucose. Antibiotics were used at the following concentrations in plates and broth: ampicillin, 100 µg/ml; spectinomycin, 100 µg/ml. Isopropyl-β-D-thiogalactopyranoside (IPTG) was used at 100 µM or at 1 mM as indicated.

Isolation and cloning of exogenous nucleic acid. *E. coli* chromosomal DNA was digested with either *Pst*I and *Hind*III or *Eco*RI and *Bam*HI and ligated into the vector pLEX5BA, cut with the same enzymes (Diederich *et al.*, Biotechniques 16:916-923 (1994)). Plasmid pLEX5BA contains: i) a Bujard promoter that has binding sites for lac repressor centered at -22 and +11 relative to the start of transcription, ii) a multiple cloning site downstream of the promoter, and iii) an *rrnBt1t2* transcriptional terminator after the multiple cloning site. Expression of fragments cloned downstream of the Bujard promoter can be induced with IPTG.

The double digestions were chosen to give fragments with a median length of 2 - 3 kb (Churchill *et al.*, 1990). The ligation mix was transformed into *E. coli* DH5α and transformants were selected on plates containing ampicillin. Colonies that grew on ampicillin were subsequently replica plated by physical transfer to a second ampicillin plate containing the inducing agent IPTG at a concentration of 100 µM. Colonies that did not grow in the presence of IPTG were chosen for further characterization.

Inhibition of bacterial growth after IPTG induction. Growth curves were carried out by back diluting cultures 1:20 into fresh media with or without 1 mM IPTG and measuring the

OD₄₅₀ every 30 minutes (min). To study transcriptional induction in exponential phase, cultures were diluted 1:10⁶ or as indicated. The viability (CFU/ml) was determined over time by plating aliquots of the culture on agar with or without IPTG.

5 There were 53 clones whose proliferation was inhibited after IPTG induction, out of approximately 50,000 *Pst*I - *Hind*III clones screened. Out of 16,000 *Eco*RI - *Bam*HI cloned fragments, the growth of 15 transformants was inhibited after IPTG induction. Thus, approximately 0.1% of the inserts resulted in IPTG sensitivity when expressed from the Bujard promoter.

Plasmids pJB1 through pJB68 were cut with restriction enzymes to release the cloned inserts. Sizes ranged from 450 bp to 6 KB. Clones that had inhibition of colony formation after induction with 100 mM IPTG and had inserts less than 1050 bp were selected for further study. Certain plasmid derivatives and strains are listed in Table 1.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187

they coded for an RNA fragment that was identical to a portion of the wild type mRNA from a chromosomal locus. About 1/13,000 clones produced an inhibitory antisense RNA.

Table 1. *E. coli* K-12 strains (all F⁻) and plasmids

Strain	Genotype/phenotype
DH5 α	supE44 Δ lacU169(ϕ 80lacZ Δ M15) <i>hsdR17 recA1 endA1 gyrA96 thi-1 relA1</i>
JRG58201	Δ (<i>nadC-ampD-ampE-aroP</i>) <i>ampG::kan</i>
Plasmids	Vector/Insert
pLEX5BA	Ap ^r cloning vector
pJB3	pLEX5BA with an 836 bp <i>Pst</i> I/ <i>Hind</i> III fragment of <i>secA</i> in sense orientation
pJB37	pLEX5BA with a 546 bp <i>Pst</i> I/ <i>Hind</i> III fragment of <i>lepB</i> in antisense orientation
pJB53	pLEX5BA with a 714 bp <i>Pst</i> I/ <i>Hind</i> III fragment of <i>viaA</i> in antisense orientation
pJB57	pLEX5BA with a 1050 bp <i>Eco</i> RI/ <i>Bam</i> HI fragment of <i>ddlB</i> in antisense orientation
pJB58	pLEX5BA with a 783 bp <i>Eco</i> RI/ <i>Bam</i> HI fragment with homology to <i>fimF</i> and <i>fimD</i>
pJB59	pLEX5BA with a 451 bp <i>Eco</i> RI/ <i>Bam</i> HI fragment of <i>orf1/ampG</i> in antisense orientation
pJB60	pLEX5BA with a 619 bp <i>Eco</i> RI/ <i>Bam</i> HI fragment of <i>ugpB</i> in sense orientation
pCW37	pLEX5BA with a 546 bp <i>Pst</i> I/ <i>Hind</i> III fragment of <i>lepB</i> in sense orientation
pCW53	pLEX5BA with a 714 bp <i>Pst</i> I/ <i>Hind</i> III fragment of <i>viaA</i> in sense orientation
pCW57	pLEX5BA with a 1050 bp <i>Eco</i> RI/ <i>Bam</i> HI fragment of <i>ddlB</i> in sense orientation
pCW59	pLEX5BA with a 451 bp <i>Eco</i> RI/ <i>Bam</i> HI fragment of <i>orf1/ampG</i> in sense orientation
pAF Ω	Sp ^r ,St ^r derivative of pLEX5BA containing 2 kb <i>Sma</i> I Ω fragment cloned into <i>Cla</i> I site
pAF37	pAF Ω with a 546 bp <i>Pst</i> I/ <i>Hind</i> III fragment of <i>lepB</i> in antisense orientation
pAF53	pAF Ω with a 714 bp <i>Pst</i> I/ <i>Hind</i> III fragment of <i>ddlB</i> in antisense orientation
pAF57	pAF Ω with a 1050 bp <i>Eco</i> RI/ <i>Bam</i> HI fragment of <i>viaA</i> in antisense orientation
pAF59	pAF Ω with a 451 bp <i>Eco</i> RI/ <i>Bam</i> HI fragment of <i>orf1/ampG</i> in antisense orientation

Example 2

Identification of Proliferation Genes by Antisense Inhibition

Seven clones obtained as described in Example 1 inhibited bacterial proliferation and
5 contained gene fragments in an antisense orientation.

Three of the clones contained fragments of the *lepB* gene; pJB12, pJB37 and pJB40. pJB37 contains a 546 bp *HindIII* - *PstI* insert that expresses an RNA complementary to the carboxy terminal half of the *lepB* mRNA. The RNA sequence is shown in Figure 1. *LepB* encodes the leader peptidase responsible for proteolytic cleavage of the signal peptide from preproteins. Leader peptidase is required for viability and a temperature sensitive *lepB* mutant
10 lyses at the nonpermissive temperature.

A fragment of an uncharacterized ORF was cloned twice in antisense orientation in pJB39 and pJB53. The antisense clone pJB53 contains a 714 bp *PstI* - *HindIII* fragment of this putative ORF, designated herein as *viaA* (viability inhibited by antisense) and expresses an RNA
15 that is complementary to the putative sense RNA. The sequence of the antisense RNA is shown in Figure 2 and the sequence of the DNA insert is shown in Figure 3. Plasmid pJB53 has been deposited with the American Type Culture Collection (ATCC), Rockville, Maryland as accession number ____.

One antisense clone, pJB57, contained a fragment of the *ddlB* gene. The D-alanyl:D-
20 alanine ligase antisense clone, pJB57, contains a 1050 bp *EcoRI* - *BamHI* fragment (with an internal *EcoRI* site) that expresses an RNA antisense to the carboxy terminal portion of *murC* and the first 600 bases of *ddlB* including its start codon. The RNA sequence is shown in Figure 4. The product of the *ddlB* gene, the D-alanyl:D-alanine ligase, ligates 2 D-alanines together

which are destined to form part of the peptidoglycan layer. A temperature sensitive *ddlB* mutant could only grow for 40 min at the non-permissive temperature before lysing, presumably due to the compromised peptidoglycan layer. Lugtenberg E. and van Schijndel-van Dam, A., *J. Bact.* 113:96-104 (1973). There is a naturally occurring inhibitor of D-alanyl:D-alanine ligase activity, the antibiotic cycloserine.

The *ampG* operon antisense clone pJB59 contains a 451 bp *EcoRI* - *BamHI* fragment which expresses an RNA antisense to a polycistronic mRNA containing *orf1* and *ampG*. The antisense RNA ends 64 nucleotides upstream of the *ampG* coding region. The *orf1* gene product has no known function but has been thought to be dispensable. *AmpG* forms a cytoplasmic membrane pore through which peptidoglycan subunits are translocated to signal inducible β -lactamase promoters when peptidoglycan damage occurs from β -lactam antibiotics. Mutations in *ampG* cause a deficiency in recycling of the peptidoglycan layer and lose 40% of their peptidoglycan per generation but remain viable. The sequence of the antisense RNA is shown in Figure 5.

Antisense RNAs to *lepB*, *ddlB*, *viaA* and *orf1* do not encode consensus ribosome loading sites. To determine if the chromosomal inserts within pJB37, pJB53, pJB57, and pJB59 could be translated into peptides, they were analyzed by computer for ribosome loading sites and open reading frames.

Plasmids pJB37, pJB53 and pJB57 do not have significant open reading frames or consensus ribosome loading sites relative to the inducible Bujard promoter. pJB59 does have an open reading frame relative to the Bujard promoter; however, no consensus ribosome loading site occurs in the antisense reading frames. The insert in pJB57 was shown to have a promoter which

was active in an *in vitro* transcription/translation assay.

The antisense inserts in pJB37, pJB53, pJB57 and pJB59 do not encode lethal peptides. Sense plasmids were constructed to examine whether the deleterious effects on cell growth and viability could be caused by peptides produced by internal promoters in the inserts present in pJB37, pJB53, pJB57 and pJB59.

Sense orientation plasmids were generated by cutting antisense plasmids pJB57 and pJB59 with the enzymes *Bst*BI and *Acc*I which generates compatible ends. Each fragment mixture was then religated in the presence of *Acc*I, which would recleave the plasmid if it religated in the "antisense" orientation but would not recognize the altered sites created by the fragment ligating in the "sense" orientation. Plasmids pCW57 and pCW59 contained the inserts from pJB57 and pJB59, respectively, in sense orientation.

To create pCW37 or pCW53, the inserts of pJB37 and pJB53 were amplified by PCR using the sequencing primers listed above. The PCR product was cut with *Hind*III and *Sma*I, filled in with Klenow and then cloned into the *Sma*I site of pLEX5BA. pCW37 and pCW53 contain the inserts from pJB37 and pJB53, respectively, in sense orientation. The orientation of all recombinants was verified by restriction analysis.

Cell proliferation for each strain was measured by diluting overnight cultures 1:10⁶ into M9 medium. After growth to an OD₄₅₀ of 0.2, aliquots of each culture were plated on M9 agar without IPTG or M9 agar containing 1 mM IPTG. The results are shown in Figure 6. Sense plasmids pCW37, pCW53, pCW57, and pCW59 had no measurable effect on cell growth or colony formation when induced with IPTG. In contrast, cells carrying pJB37 or pJB53 were unable to form colonies when plated with IPTG. Colony formation by cells carrying pJB57 was

inhibited more than 100 fold. The colony formation of strains carrying pJB59 was inhibited more than 1000 fold. These results indicate that the inserts in pJB37, pJB53, pJB57 and pJB59 do not contain internal promoters expressing a lethal peptide in the sense orientation.

5 To test whether chromosomal inserts expressed proteins, 4 μ g of CsCl purified plasmid templates was used in an *in vitro* transcription translation system for supercoiled templates (Promega, Madison, Wisconsin). The manufacturer's instructions were followed using 10 μ Ci 35 S-methionine and 100 μ M IPTG in a 50 μ l reaction. Samples were precipitated with acetone and electrophoresed on 15% SDS PAGE. Bands were detected by exposure to a Molecular Dynamics phosphorimager and to film. No detectable peptide product was made from clones pJB37 or pJB59.

10
11
12
13
14
15
16
17
18
19
20
Kinetics of inhibition during exponential growth by antisense RNA. The kinetics of antisense inhibition was examined for strains carrying pJB37, pJB53, pJB57 or pJB59. Overnight cultures of each strain were diluted 1:20 into LB media containing 1 mM IPTG or LB media without IPTG. The OD₄₅₀ was then measured on the liquid cultures. Figure 7 shows the increase in OD₄₅₀ during exponential growth for pJB37, pJB57 and pJB59, with and without induction of antisense RNA. As seen in Figure 7, transcriptional induction of antisense RNA did not have a significant effect on the increase in cell mass until about 3 hours after induction. Similar results were obtained for a strain carrying pJB53. After induction of antisense RNA expression, a few generations of cell division presumably are required in order to dilute endogenous gene product present at the initiation of induction. The cell mass for the strains expressing the *lepB* or *viaA* antisense RNAs (pJB37 and pJB53, respectively) began to decrease by about 4 hours, suggesting that cells had begun to lyse. Microscopic examination revealed that

cells expressing *lepB* antisense RNA did lyse, whereas cells expressing *viaA* antisense RNA did not. The cell mass for strains carrying *ddlB* (pJB57) and *orfI* (pJB59) stopped increasing after about 3 hours but did not decrease, suggesting that these cells did not lyse.

5 The OD₄₅₀ of strains carrying the sense constructs described above continued to increase, both in the presence and in the absence of IPTG, showing that induction of the sense RNA had no effect on bacterial proliferation.

Inhibition of viability by antisense RNA. The viability over time of the four strains expressing the antisense RNAs described above was also examined. Overnight cultures of each strain were diluted 1:100 into LB medium containing 1 mM IPTG. Aliquots of each culture were plated on LB agar containing ampicillin at intervals thereafter to determine CFUs/ml.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
10185

IPTG. At the time of induction and every 2 hours thereafter, aliquots were plated on LB/ampicillin agar to determine CFUs/ml.

The results shown in Figure 9 indicate that the anti-proliferative effect of antisense to *lepB* or *viaA* is near maximal at 500 μ M IPTG. Cell death occurred within 2 hours. A 2 fold increase to 1000 μ M IPTG had little effect on the time of cell death. A 5 fold decrease in IPTG (100 μ M) resulted in a longer period of survival (4 hours or more) before CFU began to decrease. However, cells did not survive overnight induction at 1000 μ M IPTG, as shown in Figure 6. Very low levels of IPTG (10 μ M) had minimal effect on colony forming ability.

***lepB* antisense induction inhibits leader peptidase activity.** The *lepB* gene encodes a leader peptidase that cleaves the leader sequence from preproteins such as maltose binding protein (MBP). Endogenous leader peptidase activity was examined in the presence of pJB37 antisense RNA, by measuring processing of pre-MBP (43.38 KD) to the mature 40.7 KD protein.

An overnight culture of pJB37/DH5 α was diluted 1:100 in M9 media containing 0.2% casamino acids, 0.2% maltose, 100 μ g/ml ampicillin. The diluted culture was split into 2 flasks, one containing 1 mM IPTG and the other containing no IPTG. Samples were taken at the time of the split and at hourly intervals thereafter. Chloramphenicol was added to each sample to a final concentration of 150 μ g/ml. Samples were placed on ice, centrifuged, washed with M9 salts, and precipitated with cold 10% trichloroacetic acid (TCA). After centrifugation, the TCA precipitated material was resuspended in buffer (0.125 M Tris-HCL, pH 6.8, 0.1 M NaOH, 1% w/v SDS, 0.1% Phenol Red). The amount of protein in each precipitate was normalized to the different optical densities of the samples by using 0.1 ml of buffer per ml of culture having an OD₄₅₀ of 1.0. About 8 μ l of each resuspended precipitate was mixed with an equal volume of 2X

cracking buffer (100 mM Tris-Cl pH6.8, 200 mM DTT, 4% SDS, 0.2% bromophenol blue, 20% glycerol) and boiled for 10 minutes. Samples were electrophoresed on a 10% polyacrylamide-SDS gel as described in Sambrook *et al.*, *supra* (1989).

5 Gel-separated proteins were electrophoretically transferred to a Protran nitrocellulose membrane (Schleicher & Schuell) and the membrane incubated with rabbit anti-MBP antibody diluted to 1:10,000. After washing, the membrane was incubated with a 1:10,000 dilution of goat anti-rabbit antibody conjugated to horseradish peroxidase (BioRad). Proteins were detected with a chemiluminescence kit (DuPont-NEN) and captured on autoradiographic film. The image
10 was scanned using a Molecular Dynamics densitometer. The results showed that unprocessed MBP is detectable after induction of *lepB* antisense transcription with IPTG. No unprocessed MBP is detectable in untreated cells. The data demonstrate that *lepB* antisense RNA results in inhibition of the endogenous bacterial gene product.

Lethality of pJB59 antisense RNA requires inhibition of *orf1*. The polycistronic mRNA that encodes *ampG* in *E. coli* has an upstream ORF, termed *orf1* herein. The antisense RNA made by pJB59 is complementary to *orf1*. In order to determine whether translational inhibition of *orf1* or of *ampG* was the cause of the lethality of pJB59 antisense RNA, pJB59 was transformed into the *ampG* null strain, JRG58201. Jacobs, C., *et al.*, EMBO J. 13:4684-4694 (1993). JRG58201 has a defect in the *ampG* gene but has no known defect in the *orf1* gene.

20 Growth and viability of JRG58201 containing pJB59 was inhibited after IPTG induction. Thus, the anti-proliferative effect of the antisense RNA of pJB59 requires at least the inhibition of *orf1* gene expression. Whether the anti-proliferative effect of the *ampG* gene product is also involved cannot be determined from this experiment.

Effect of different antibiotic selectable markers on antisense inhibition of growth.

The possibility that IPTG induction was sensitizing the cells to ampicillin was examined by transferring the inserts from plasmids pJB37, pJB53, pJB57 or pJB59 to the vector pAF Ω .

- 5 Plasmid pAF Ω confers resistance to ampicillin, spectinomycin, and streptomycin. The resulting constructs were then tested for inhibition of bacterial growth using spectinomycin as the selectable marker rather than ampicillin.

pAF Ω was constructed by isolating the *Sma*I fragment containing Omega (Sp^r , Str^r) of pHP45. Prentki and Kirsch, Gene 29:303-313 (1984). The omega fragment was then ligated into the filled-in *Cla*I site of pLEX5BA.

To generate pAF37, pAF53, pAF57, and pAF59, the inserts of pJB37, pJB53, pJB57, pJB59, respectively, were amplified by PCR. The PCR products of pJB37 and pJB53 were cut with *Hind*III, filled in with Klenow, and cut with *Eco*RI before ligation into *Eco*RI-*Sma*I cut pAF Ω . The PCR products of pJB57 and pJB59 were cut with *Pst*I and *Eco*RI and ligated into identically cut pAF Ω . Ligation reactions were transformed into DH5 α and ampicillin spectinomycin resistant transformants were selected. Generation of the correct constructs was verified by restriction analysis.

20 Overnight cultures of strains carrying plasmids pAF37, pAF57 or pAF59 were diluted 1:100 in LB medium containing spectinomycin. After growth to an OD₄₅₀ of 0.2, cells were plated on ampicillin, ampicillin with 1 mM IPTG, spectinomycin, or spectinomycin with 1 mM IPTG, and CFUs/ml of culture were determined. The results are shown in Figure 10.

Figure 10 shows that induction of antisense nucleic acid under spectinomycin selection inhibits bacterial proliferation to the same extent as does antisense nucleic acid under ampicillin

selection. Similar results were obtained for pAF53. These results suggest that the inhibition of bacterial proliferation observed with these constructs is not due to an effect on ampicillin resistance.

5 In order to examine the effectiveness of the antisense RNAs in a related species, *Salmonella typhimurium* was transformed with pJB37 or pJB53. The ampicillin-resistant transformants were plated on LB with or without 1 mM IPTG. Growth was greatly inhibited after induction of the antisense RNA, suggesting that sufficient complementarity existed between the antisense RNAs and *S. typhimurium* genes to inhibit production of proliferation gene products in this species as well as in *E. coli*.

10 The data described above indicate that antisense clones inhibit cell growth by producing an antisense RNA. Transcription of the cloned fragment must occur because IPTG induction is necessary to inhibit proliferation. None of the inserts are lethal in a sense orientation. Thus, any peptide possibly made by the insert in pJB57 is not responsible for lethality. No detectable peptide product is made from the antisense RNA of clones pJB37 or pJB59 *in vitro*. The pJB37 (*lepB*) antisense RNA was shown to decrease leader peptide processing activity.

Example 3

Identification of Proliferation Genes by Sense Inhibition

20 Four of the clones described in Example 1 inhibited bacterial proliferation and contained gene fragments in sense orientation. Plasmid pJB3 contained a fragment of the *secA* gene in sense orientation. Plasmid pJB60 contained a fragment of the *ugpB* gene in sense orientation. Plasmids pJB55 and pJB58 each contained a fragment in the sense orientation which expressed

portions of genes having sequence homology to the *E. coli fimF* and *fimD* genes (located at 34 min on the *E. coli* map).

The insert in the *secA* dominant lethal clone, pJB3, was identified as an 836 bp *Pst*I-
5 *Hind*III fragment coding for *yacA* and the N-terminal 39 amino acids of the *secA* protein. The DNA sequence of the insert is shown in Figure 11. *SecA* is an ATP-dependent translocase of secreted proteins located at the cytoplasmic face of the membrane. *SecA* protein regulates its own translation by binding to a site overlapping the ribosome binding site, which causes dissociation of a preformed 30S-tRNA-fMet-mRNA ternary complex. Since this site is located on the cloned fragment in pJB3, the sense RNA produced by pJB3 after induction could be binding and potentially sequestering most of the *secA* protein, resulting in a fatal deficiency in trans-membrane transport. Alternatively, the truncated *secA* peptide produced by pJB3 could be poisoning the membrane complex or binding the wild type mRNA, down regulating translation of *secA*.

10
15
An antisense RNA complementary to *secA* mRNA would also be expected to inhibit bacterial proliferation. The nucleotide sequence of such an antisense RNA is shown in Figure 12.

20 The dominant lethal *ugpB* sense clone, pJB60, contains a 619 bp *Eco*RI - *Bam*HI fragment which encodes the amino-terminal portion of the *sn*-glycerol-3-phosphate binding protein.

The insert in the dominant lethal clone, pJB58, is a *Bam*HI - *Eco*RI fragment approximately 783 bp long. The plasmid insert contains two partial ORFs in a sense orientation. One partial ORF encodes the C-terminal portion of a protein which shows 59% similarity to

FimD protein. The other partial ORF encodes the N-terminal portion of a protein with 56% similarity to FimF protein.

To the extent not already indicated, it will be understood by those of ordinary skill in the art that any one of the various specific embodiments herein described and illustrated may be further modified to incorporate features shown in other of the specific embodiments.

The foregoing detailed description has been provided for a better understanding of the invention only and no unnecessary limitation should be understood therefrom as some modifications will be apparent to those skilled in the art without deviating from the spirit and scope of the appended claims.